# Efficacy of Carbendazim and other Synthetic Fungicides on Taro Leaf Blight Disease caused by *Phythophtora Colocasiae*

*Mbadianya, J. M.* [*1, 2], *Ugwuoke, K.I.*[3], *Baiyeri, K.P.*[3], *Shi, D.*[1] *and Chen, C.*[1]

[1]*College of Plant Protection, Nanjing Agricultural University, Key Laboratory of Pesticide Science, Jiangsu Province, Nanjing 210095, China*
[2] *Department of Crop Science and Horticulture, Nnamdi Azikiwe University, Awka, 5025, Nigeria*
[3] *Department of Crop Science, University of Nigeria, Nsukka*

KEYWORDS

*Cocoyam leaf blight,*
*Fungicides,*
*Mycelia growth and resistance*

ABSTRACT

*The results of the serial concentrations of different synthetic fungicides, phytopathogenic fungi and their interactions on isolates from cocoyam had significant effects ($P \leq 0.05$) on the percentage inhibition and mycelia growth. P. colocasiae (12.50 mm) treated with hexaconazole had the lowest mycelia growth rate followed by Botryodiplodia theobromea (21.28 mm) with fludioxonil and carbendazim (24.03 mm) while Rhizopus spp (85.83 mm) and F. solani (78.798 mm) on metaxyl recorded the highest mycelia growth. The effects of different environmental stress, fungi and their interaction on mycelia growth showed significant difference on the organisms, the stress and their interactions. No mycelia growth were seen when B. theobromea were subjected to $H_2O_2$ (5.00 mm), followed by P.colocasia (14.67 mm).*

*In conclusion, fungicides found to be effective for the control of cocoyam leaf blight in order of merit were carbendazim, fludioxonil, hexaconazole and mancozeb. Metaxyl recorded no or little effect on the inhibition of mycelia growth of all the four fungi organisms isolated from cocoyam as the organisms have developed resistance to it. It should be used in mixture with other fungicides with different mode of actions for example mancozeb.*

*CORRESPONDING AUTHOR

*ji.mbadianya@unizik.edu.ng*

## INTRODUCTION

Cocoyams, *Colocasia esculenta* are important perennial food crops cultivated for their edible roots (Owusu-Darko *et al*., 2014). The crop is primarily grown in tropical and sub-tropical countries and is one of the major food crops in Nigeria (Aniekwe, 2015, Omeje *et al*., 2015).

Despite their importance and usage, their production in Nigeria are heavily affected by plant diseases especially fungi including; *Phyllosticta colocasiophila*, *Phytophthora colocasia*, *Fusarium spp*, *Phythium spp*, *Aspergillus niger*, *Sclerotia rolfsii* and *Botryodiplodia theobromae* (Anukworji *et al*., 2012; Omeje *et al*., 2015) but the most devastating fungus is *P. colocasia* which causes leaf blight. Symptoms are small, brown, water-soaked lesions on the leaves which enlarge and coalesce into large lesions with yellow exudate, defoliation and death.

The infection of taro leaf blight has reduced taro production rate by 30-40%, leading to heavy reduction in quality and quantity of corm and hindering the commercialization of taro corm product in many taro producing countries (Bandyopadhyay *et al*., 2011; Mbong *et al*., 2013). Control measures need to be adopted to reduce the effect of taro leaf blight (TLB) below its economic injury level and hence increase taro production. TLB has been controlled using cultural strategies including field sanitation, roguing, clean planting stocks, and use of resistant varieties (Aggarwal and Mehrotra, 1987; Nelson *et al*., 2011) such as *Trichoderma* species, *Rhizobacterium* species) (Chukwu and Enyiukwu, 2021). However, due to the drastic reduction in yield of taro, the use of fungicide is justified to ensure quick response.

Fungal infection has been worldwide estimated to reduce yield of crops to 20% if not treated, thus use of fungicides has become wide spread recently in Agriculture (Rohr *et al*., 2017). Fungicide controls the fungal infection during the establishment and growth of a crop, it enhances the productivity of crop and decrease defects (Bandyopadhyay *et al*., 2011). Some fungicides that can be used to control taro leaf blight disease includes single or compound synthetic fungicides such as mancozeb, diflolatan, and ridomil (metalaxyl) or copper (Aggarwal and Mehrotra, 1987; Cox and Kasimani, 1988).

To improve the efficiency of production, this fungus must be controlled and managed through the use of some synthetic fungicide benzimidazole.

Benzimidazoles is a group of fungicides commercially available as benomyl, carbendazim (MBC), thiophanate-methyl, thiabendazole and fuberidazol, known for their broad-spectrum activities against several fungal pathogens namely ascomycetes, some basidiomycetes and deuteromycetes (Leadbeater, 2014). These fungicides can be applied to cereals, fruits, vegetables and vines and also used in postharvest handling of crops (Duan *et al*., 2014; Y. Duan et al., 2019; Oliver and Hewitt, 2014) against *Cercospora spp*., *F. spp*, *Botrytis cinerea Colletochrichum spp*., powdery mildew, *Erysiphe spp* and *Oidium spp* crops (Duan *et al*., 2019; Oliver and Hewitt, 2014). Carbendazim is the most common and widely used to manage crop diseases in cereals, roots and fruits (Zhou *et al*., 2016). The combination of carbendazim and diethofencarb is a good control measure against *B. cinerea* although the appearance and spread of the two fungicides caused problems (Leroux and Fritz, 1984).

Benzimidazoles interrupt pathogen energy metabolism through their selective bindings with high affinity to pathogen β-tubulin and inhibition of microtubule polymerization resulting to destruction of cell structure and death of pathogen. They are the important inhibitors of β-tubulin polymerization against several plant pathogenic fungi such as *B. cinerea*, *Cercospora spp*., *C. spp*., *F. spp*., *E. spp* and *O. spp* (Duan *et al*., 2019). This study was therefore, to determine the efficacy of synthetic fungicide carbendazim on the leaf blight of cocoyam caused by *P. colocasiae* with the following objectives to:

(a) Determine the sensitivity and resistance of *P. colocasiae* to carbendazim.
(b) Assess the effect of carbendazim and other fungicides on other fungi organisms associated with cocoyam
(c) Evaluate the effect of *P. colocasiae* and other fungi organisms isolated from cocoyam on environmental stress

## MATERIALS AND METHODS

### Experiment sites

Laboratory experiments were conducted at the Department of Crop Science, Teaching and Research Farm, Faculty of Agriculture, University of Nigeria, Nsukka.

### Collection of P. colocasiae isolates

Cocoyam leaves with leaf blight symptoms were collected from Department of Crop Science research farm, University of Nigeria, Nsukka. The leaves were disinfected with 0.5% (vol/vol) of Naclo for 1 minute and 75% (vol/vol) ethanol for 30s, rinsed three times with distilled water and cultured in potato dextrose agar (PDA) plates PDA which contains 200g of boiled potato tubers, 20g of dextrose and 15g of agar with distilled water per litre, amended with streptomycin sulfate (98% a.i., 50 μg/ml; Solarbio Science & Technology Co., Ltd., Beijing, China) and incubated for 3 days. Mycelia growth emerging were cut from the colony margins and transferred to fresh PDA plates. Pure cultures of the isolates were collected by sub-culturing on PDA under 12/12h light/darkness photoperiod using near-ultraviolet (NUV) light at 25 $^0$C. a single conidium of the isolates obtained were maintained PDA slant at 4 $^0$C for further use.

### Fungicides and media

All fungicides of pure technical grade were used for the experiment. Fludioxonil (97.9% a.i.), carbendazim (98% a.i), metalxyl, mancozeb and hexaconazole (98.7% a.i.) were dissolved in methanol to obtain $10^4$ (μg) $ML^{-1}$ of the stock solution were preserved in the dark at 4 $^0$C. PDA were prepared with 200g of potato, 20 g of glucose and 16 g of agar $L^{-1}$ of distilled water where Potato dextrose broth (PDB) used for these experiments also has the same composition but lacks agar.

### Evaluation of sensitivity of *P. colocasia* to carbendazim

The sensitivity and resistance test of *P. colocasia* and other fungi organisms isolated from cocoyam were estimated using mycelia growth assay. The other fungi organisms were *Fusarium solani, Rhizopus spp* and *Botryodioplodia theobromea*.

Mycelia plugs (5-mm in diameter) were prepared from one-third outside of the active margin of colony, transferred to new PDA plates and used for the experiment treated with carbendazim, mancozeb, metalaxyl, fludioxonil and hexaconazole at the serial concentrations of 0, 10, 20, 40, 80 and 100 μg/ml and incubated at 25 0C for 6 days. The diameters were investigated when the colonies without fungicide approached to eight cm. Data on mycelia growth in terms of colony diameter were obtained by averaging the diameters at

perpendicular directions from which 5-mm in diameter were subtracted. The percentages of growth inhibition was calculated and arcsine-transformed prior to statistical analysis using the percentage transformation formula (Inhibition rate = ARSIN (SQRT (ab/100))*180/3.1415926) prior to statistical analysis. The treatments were replicated three times for each strain of the six pathogens and experiments performed two times.

**Virulence test assay**

To determine whether the organisms associated with cocoyam leaves can cause infection, virulence test assay were done on healthy cocoyam leaves.

Mycelia plugs (5mm in diameter) from a 6 days old PDA culture of each isolates were placed on the surface of the fruits over artificial wound for easy penetration of the pathogen, keep moist with a piece of moistened absorbent cotton and incubated in a growth chamber at 25 0C under 12-h photoperiod and 85% relative humidity. After 6 days, lesion diameters were measured as the mean of two diameters (cm) at perpendicular angles. The experiment was done two times with three replications

**Osmotic stress assay**

Mycelia plug (5mm in diameter) was taken from the edge of a 5 days old colony and transferred to PDA amended with 1.2 M NaCl, 1.2 M KCl, 0.5 M $CaCl_2$, 0.3 mg/ml Congo red and 32µg/ml $H_2O_2$. PDA without osmoticum served a control; each treatment was replicated three times and incubated at 25 $^0$C for seven days. The percentages of growth inhibition were calculated and arcsine-transformed prior to statistical analysis as they were not evenly distributed. The experiment was conducted two times.

**Experimental design**

The experiments were laid out as factors in a completely randomized design (CRD) with three replications. Factor A were the different plant fungi, factor B were the fungicides and factor C were different concentrations.

**Statistical analysis**

Data were analyzed using a GENSTAT statistical software package, 12.0 Release 4.23 (Payne, 2009). Colony diameters, mycelia weights and lesion areas were estimated in a completely randomized design (CRD) with three replications. Means were compared with Fisher's protected least significant difference at 5% probability level as outlined by (Obi, 2002).

**RESULTS**

**Evaluation of sensitivity of *P. colocasia* to carbendazim and other fungicides**

The results showed that serial concentrations of different synthetic fungicides, phytopathogenic fungi and their interactions had significant effect ($P \leq 0.05$) on the percentage inhibition and mycelia growth of isolates from cocoyam. Five fungicides at concentrations: 0.10, 20, 40, 80 and 100 µg/ml evaluated *in vitro* against *P. colocasiae, F. solani, R. spp* and *B. theobromea* exhibited wide range of mycelia growth and inhibition of the pathogens. *P. colocasia*e (12.50 mm) treated with hexaconazole had the lowest mycelia growth rate followed by *Botryodiplodia theobromea* (21.28 mm) with fipronil and carbendazim (24.03 mm) while *Rhizopus spp* (85.83 mm) and *F. solani* (78.798 mm) on metaxyl recorded the highest mycelia growth (Fig 1). Low mycelia growths were seen on all fungal isolates tested on concentration 100 µg/ml when compared with the untreated plates (Fig 2). Concentration 100 µg/ml of carbendazim (7.00 mm) and mancozeb (5.00 mm) recorded the lowest mycelia growth (Fig 3). The organism *B. theobromea* at concentrations 10, 20, 40, 80 and 100 µg/ml of carbendazim and 80 and 100 µg/ml of mancozeb did not grow likewise *P. colocasiae* at concentrations 10 -100 µg/ml of hexaconazole and 20 -100 µg/ml of carbendazim. Concentrations 40 – 100 µg/ml of carbedazim did not allow rhizopus to grow (Fig 4). The results revealed that all the five fungicides tested at concentrations 40 -100 µg/ml significantly inhibited mycelial growth over untreated plates. Furthermore, the percentage mycelia growth inhibition of carbendazim and mancozeb increased with increased in concentration of fungicides tested. Highest percentage inhibitions of *P. colocasiae* were observed on hexaconazle (83.33%) and carbendazim (75.97%). However, fungicides carbendazim, fipronil, hexaconazole and mancozeb were considered to be good for the inhibition of *B. theobromea* and *P. colocasiae*. The effects of different environmental stress, fungi and their interaction on mycelia growth showed significant difference on the organisms, the stress and their interactions. No mycelia growth were seen when *B. theobromea* were subjected to $H_2O_2$ (5.00 mm), followed by *P.colocasia* (14.67 mm) (Fig 5).

**Table 1:** Mycelia growth rate and virulence test assay of *P. colocasia , B.theobromea, R. spp* and *solani*

| Fungi | Virulence assay | mycelia weight |
|---|---|---|
| *Rhizopus spp* | 8.5 b | 0.023 |
| *B. theobromea* | 10 b | 0.104 |
| *F. solani* | 19.17 a | 0.062 |
| *P. colocasia* | 10 b | 0.453 |
| F-LSD $_{(0.05)}$ | 3.863 | NS |




**Fig 1**: Effects of different synthetic fungicides and their concentration on the percentage inhibition of mycelia growth of *Phytophthora colocasiae.*




**Fig 2:** Effects of different synthetic fungicides and their concentration on the percentage inhibition of mycelia growth of *Fusarium solani***.**




**Fig 3**: Effects of different synthetic fungicides and their concentration on the percentage inhibition of mycelia growth of *Botryodiploma theobromea.*




**Fig 4:** Effects of different synthetic fungicides and their concentration on the percentage inhibition of mycelia growth of *Rhizopus spp.*




**Fig 5:** Effects of different environmental stress on the percentage inhibition of mycelia growth of different fungi organisms isolated from cocoyam.

DISCUSSION

The results revealed that all the five fungicides tested at concentrations 40 -100 µg/ml significantly inhibited mycelial growth over untreated plates. Furthermore, the percentage mycelia growth inhibition of carbendazim and mancozeb increased with increased in concentration of fungicides tested. Highest percentage inhibitions of P. colocasiae were observed on hexaconazle (83.33%) and carbendazim (75.97%). However, fungicides carbendazim, hexaconazole , mancozeb, and fludioxonil were considered to be more effective in the inhibition of *B. theobromea* and *P. colocasiae.*

Early research on the mechanism of action of carbedazim dwell on DNA and RNA synthesis which were described as secondary and primary effects respectively (Davidse, 1995;Davidse, 1986). In arresting nuclear division of fungi, carbendazim fungicide gives a striking resemblance to secondary plant metabolite colchicine, which disrupt mitosis and meiosis in animal and plant cells by inactivating the spindle (Zhou *et al*., 2016). The biochemical analysis of mechanism of action of carbendazim indicate the antimitotic activity of carbendazim in fungi facilitated through binding to fungal tubulin (Davidse, 1995; Zhou *et al*., 2016). They are greatly important in controlling many plant pathogenic diseases of crops. Carbendazim is a strong inhibitor of tubulin polymerization, antifungal in action and interfere with parasites energy metabolism through their selective bindings with high affinity to pathogen β-tubulin resulting to destruction of cell structure and death of pathogen (Chen *et al*., 2009; Lacey, 1990; Prichard, 1970a, 1970b; Schmit, 2013; Tejada *et al*., 1987; Y. Zhou *et al*., 2016). Hexaconazole is used widely for the management of anthracnose, powdery mildew, late blight, early blight, downy mildew and grey mildew diseases on crops due to its broad spectrum activity (Zhou *et al*., 2017).

Metalaxyl spectrum of activity broadened to control more diseases when it is used with other fungicides such as dithiocarbamate, phythalimides or copper fungicides and at the same time thus buildup of resistant fungal strains may be delayed and prevented (Sukul and Spiteller, 2000; Yao *et al*., 2009). Metalaxyl inhibits ribosomal RNA synthesis through RNA polymerization by acting on the polymerase 1 complex of rRNA synthesis which is the target site (Agrios, 2005). Resistance isolates of metalaxyl were detected in *Pseudoperenospora cubensis, P. infestans, P. tabacina* and *P. viticola*, therefore it was recommended that it should use in mixture with other fungicides with different mode of actions for example mancozeb (Agrios, 2005)

The effects of different environmental stress, fungi and their interaction on mycelia growth showed significant difference on the organisms, the stress and their interactions. No mycelia growth were seen when *B. theobromea* were subjected to $H_2O_2$ (5.00 mm), followed by *P.colocasia* (14.67 mm).

In conclusion, fungicides found to be effective for the control of cocoyam leaf blight in order of merit were carbendazim, fludioxonil, hexaconazole and mancozeb. Metaxyl recorded no or little effect on the inhibition of mycelia growth of all the four fungi organisms isolated from cocoyam as the organisms have developed resistance to it. It should be used in mixture with other fungicides with different mode of actions for example mancozeb.

**REFERENCES**


Aggarwal, A. and Mehrotra, R. (1987). Control of Phytophthora leaf blight of taro (Colocasia esculenta) by fungicides and roguing. Phytoparasitica, 15, 299-305.

Agrios, G. N. (2005). Plant pathology: Elsevier.

Aniekwe, L. (2015). Improving food security with cocoyam production by the smallholder female farmers in Ebonyi State, Southeastern Nigeria. Journal of Biology, Agriculture and Healthcare, 5(13), 15-21.

Anukworji, C., Putheti, R. R. and Okigbo, R. (2012). Isolation of fungi causing rot of cocoyam (Colocasia esculenta (L.) Schott) and control with plant extracts:(Allium sativum, L., Garcinia kola, heckel., Azadirachta indica, L. and Carica papaya, L.). Global Advanced Research Journal of Agricultural Science, 1(1), 1-15.

Bandyopadhyay, R., Sharma, K., Onyeka, T., Aregbesola, A. and Kumar, P. L. (2011). First Report of Taro (Colocasia esculenta) Leaf Blight Caused by Phytophthora colocasiae in Nigeria. Plant disease, 95(5), 618.

Chen, C.-J., Yu, J.-J., Bi, C.-W., Zhang, Y.-N., Xu, J.-Q., Wang, J.-X. and Zhou, M.-G. (2009). Mutations in a β-tubulin confer resistance of Gibberella zeae to benzimidazole fungicides. Phytopathology, 99(12), 1403-1411.

Chukwu, L. A. and Enyiukwu, D. N. (2021). Direct Research Journal of Biology and Biotechnology.

Cox, P., and Kasimani, C. (1988). Control of taro leaf blight using metalaxyl. International Journal of Pest Management, 34(1), 81-84.

Davidse, L. (1995). Biochemical and molecular aspects of the mechanisms of action of benzimidazoles, N-phenylcarbamates and N-phenylformamidoximes and the mechanisms of resistance to these compounds in fungi. Modern selective fungicides.

Davidse, L. C. (1986). Benzimidazole fungicides: mechanism of action and biological impact. Annual review of phytopathology, 24(1), 43-65.

Duan, Y.-B., Ge, C.-Y., and Zhou, M.-G. (2014). Molecular and biochemical characterization of Sclerotinia sclerotiorum laboratory mutants resistant to dicarboximide and phenylpyrrole fungicides. Journal of Pest Science, 87(1), 221-230.

Duan, Y., Xin, W., Lu, F., Li, T., Li, M., Wu, J., . . . Zhou, M. (2019). Benzimidazole-and QoI-resistance in Corynespora cassiicola populations from greenhouse-cultivated cucumber: An emerging problem in China. Pesticide biochemistry and physiology, 153, 95-105.

Lacey, E. (1990). Mode of action of benzimidazoles. Parasitology Today, 6(4), 112-115.

Leadbeater, A. (2014). Plant health management: Fungicides and antibiotics.

Leroux, P. and Fritz, R. (1984). Antifungal activity of dicarboximides and aromatic hydrocarbons and resistance to these fungicides. Paper presented at the Symposium series-British Mycological Society.

Mbong, G., Fokunang, C., Fontem, L., Bambot, M. and Tembe, E. (2013). An overview of Phytophthora colocasiae of cocoyams: A potential economic disease of food security in Cameroon. Discourse Journal of Agriculture and Food Sciences, 1(9), 140-145.

Nelson, S., Brooks, F. and Teves, G. (2011). Taro leaf blight in Hawaii.

Obi, I. (2002). Statistical methods of detecting differences between treatment means and research methodology issues in laboratory and field experiments. Nsukka, Nigeria.

Oliver, R. P. and Hewitt, H. G. (2014). Fungicides in crop protection: Cabi.

Omeje, T., Ugwuoke, K., Aba, S., Eze, S., Ogwulumba, S., and Ezema, R. (2015). Field management of Phytophthora blight disease of cocoyam Colocasia esculenta L.) With spray regimes of selected fungicides in Nsukka, south eastern Nigeria. Agro-Science, 14(2), 36-45.

Owusu-Darko, P. G., Paterson, A. and Omenyo, E. L. (2014). Cocoyam (corms and cormels)—An underexploited food and feed resource. Journal of Agricultural Chemistry and Environment, 3(01), 22.

Payne, R. W. (2009). GenStat. Wiley Interdisciplinary Reviews: Computational Statistics, 1(2), 255-258.

Prichard, R. (1970a). Mode of action of the anthelminthic thiabendazole in Haemonchus contortus. Nature, 228(5272), 684-685.

Prichard, R. (1970b). Mode of action of the anthelminthic thiabendazole in Haemonchus contortus. Nature, 228, 684-685.

Rohr, J. R., Brown, J., Battaglin, W. A., McMahon, T. A. and Relyea, R. A. (2017). A pesticide paradox: fungicides indirectly increase fungal infections. Ecological Applications, 27(8), 2290-2302.

Schmit, J. (2013). In vitro anti-cancer effects of benzimidazoles on the canine osteosarcoma D17 cell line.

Sukul, P. and Spiteller, M. (2000). Metalaxyl: persistence, degradation, metabolism, and analytical methods. Reviews of environmental contamination and toxicology, 164, 1-26.

Tejada, P., Sanchez-Moreno, M., Monteoliva, M. and Gomez-Banqueri, H. (1987). Inhibition of malate dehydrogenase enzymes by benzimidazole anthelmintics. Veterinary parasitology, 24(3-4), 269-274.

Uguru, M., Baiyeri, K. and Aba, S. (2011). Indicators of climate change in the derived savannah Niche of Nsukka, South-Eastern Nigeria. Agro-Science, 10(1).

Yao, K., Zhu, L., Duan, Z., Chen, Z., Li, Y. and Zhu, X. (2009). Comparison of R-metalaxyl and rac-metalaxyl in acute, chronic, and sublethal effect on aquatic organisms: Daphnia magna, Scenedesmus quadricanda, and Danio rerio. Environmental Toxicology: An International Journal, 24(2), 148-156.

Zhou, J., Li, J., Li, J., Ren, G., Ren, Y. and Ma, H. (2017). Structure and enhanced antifungal activity of a divalent cobalt (II) complex with hexaconazole. Chemical Research in Chinese Universities, 33(6), 864-868.

Zhou, Y., Xu, J., Zhu, Y., Duan, Y. and Zhou, M. (2016). Mechanism of action of the benzimidazole fungicide on Fusarium graminearum: Interfering with polymerization of monomeric tubulin but not polymerized microtubule. Phytopathology, 106(8), 807-813.